



## International Journal of Advanced Psychiatric Nursing

E-ISSN: 2664-1356  
P-ISSN: 2664-1348  
www.psychiatricjournal.net  
IJAPN 2020; 2(2): 21-25  
Received: 24-05-2020  
Accepted: 27-06-2020

**Zulkiflu Musa Argungu**  
Department of Nursing  
Sciences, Faculty of Allied  
Health, Usman Danfodiyo  
University, Sokoto, Nigeria

**Ado Shehu**  
Department of Nursing  
Sciences, Maryam Abacha  
American University of Niger  
Maradi, Niger

**Deborah JY**  
Asokoro District Hospital,  
Julius Nyerere Street Asokoro  
Abuja, Nigeria

**Jama MD**  
Maitama General Hospital  
Abuja, Nigeria

**Corresponding Author:**  
**Zulkiflu Musa Argungu**  
Department of Nursing  
Sciences, Faculty of Allied  
Health, Usman Danfodiyo  
University, Sokoto, Nigeria

### Prevalence of and risk factors for depressive symptoms among people living with HIV/AIDS receiving antiretroviral treatment in Nigeria

**Zulkiflu Musa Argungu, Ado Shehu, Deborah JY and Jama MD**

#### Abstract

**Background:** Depression is a common mental disorder (CMD) with significant contributions to the burden of disease. It can lead to high social, economic and individual costs because it accounts for one-third of the days missed at work and a fifth of all primary health-care appointment.

**Objective:** We aimed to explore the prevalence of and risk factors for depressive symptoms among people living with HIV/AIDS (PLWHA) receiving antiretroviral treatment (ART) in Kebbi State, Nigeria.

**Method:** A cross-sectional study evaluating adult PLWHA receiving ART in three designated clinical hospitals was conducted. The validated Beck Depression Inventory (BDI) was used to assess depressive symptoms, ineligible participants.

**Result:** Multinomial regression analysis was used to explore the risk factors for depressive symptoms. 348 participants were finally included in all analyses. 40.3% were found to have depressive symptoms with 13.7% having mild depressive symptoms and 26.6% having moderate to severe depressive symptoms.

**Conclusion:** The results of multinomial regression analysis suggested that being married or living with a partner, recent experience of ART-related side effects, and/or history of HCV infection were positively associated with mild depressive symptoms, while increasing age was positively associated with moderate to severe depressive symptoms.

**Keywords:** Prevalence, risk factor, depressive symptoms, HIV/AIDS, antiretroviral treatment

#### Introduction

Depressive disorder is one of the common mental disorders (CMDs) with significant contributions to the burden of disease and disability in low- and middle-income countries<sup>[1]</sup>. It is believed that depression can lead to high social, economic and individual costs because they account for one-third of the days missed at work and a fifth of all primary health-care appointments<sup>[2]</sup>. CMDs have been known to increase disease progression among PLHIV and are the leading cause of disability among PLWHA<sup>[3]</sup>. Some of the reasons proffered for the higher prevalence of CMDs among PLWHA include decreased adherence to antiretroviral therapy<sup>[4]</sup>, emotional stress associated with the knowledge of one's HIV-positive status, externalized and internalized AIDS-related stigma and a compromised immune system<sup>[6]</sup>.

A systematic review of PLWHA depression on anti-retroviral therapy (ART) in sub-Saharan Africa reported a pooled estimate of 31.2 percent (95% CI 25.5–38.2%) of the prevalence of depressive symptoms in this population. This systematic review included 30 studies from 10 countries in sub-Saharan Africa (three studies were from Nigeria) with a combined sample size of more than 10,000 participants<sup>[7]</sup>. In a survey in Lagos, Nigeria, about 29 percent of PLHIV was diagnosed with depression using the Hamilton Depression Scale Rating,<sup>[8]</sup> although this survey had a low sample size (n = 87). Depression is linked to poor health-related quality of life and poor adherence to medication regimens among PLHIV<sup>[9, 10]</sup>. The prevalence of depression was found to be approximately 40 per cent in a survey conducted in Kano State, Nigeria, with 162 PLWHA aged 15 to 25 years using the Hospital Anxiety and Depression Scale<sup>[11]</sup>. Another study involving 310 PLHIV in the town of Zaria, Kaduna State, using the Center for Epidemiological Studies Depression Scale (CES-D), found that 21 percent of PLWHA had severe depressive symptoms<sup>[12]</sup>. The comparatively large variability in the findings of studies examining the incidence of PLWHA depression in Nigeria can be partly due to the relatively limited sample sizes and variations in the geographical location where these studies have been performed.

Sub-Saharan Africa accounts for about 70 % of the global HIV burden, with a prevalence of 4.4% as of 2015 [13]. Nigeria is the most populous country in Africa with an estimated population of more than 170 million people [14]. HIV prevalence in Nigeria as of 2015 was 3.2% (about 3.5 million people living with HIV in the country) [15]. The rise of the HIV/AIDS pandemic has brought with it many pressures on society, the health care system, families and individuals who are diagnosed with HIV [16]. Throughout the start of the pandemic, initial efforts centered primarily on how to minimize the levels of infection and mortality of those already infected with the virus [16]. More and more attention is being paid internationally to improving the quality of life of people living with HIV/AIDS [17]. This trend is particularly important given the positive steps that have already been taken in the fight against AIDS over the years with regard to improved anti-retroviral drugs (ARVs), leading to an increase in the life expectancy of people infected with the virus, as well as a decrease in the rate of new infections [17].

Despite the availability and expanded use of ART, lifelong adherence to treatment and persistent HIV infection may result in psychological problems in PLWHA [18]. A recent systematic review in Asia concluded that the prevalence of depressive symptoms among PLWHA was greater than 60% across research settings [19], and much higher than that identified in the general population [20]. Untreated depressive symptoms has been found to be associated with poor ART adherence and rapid progression to AIDS [21, 22], while the alleviation of depressive symptoms has been evidenced to benefit both ART adherence and clinic attendance [23]. Hence, knowledge of risk factors for depressive symptoms is necessary to be elucidated for clinicians for better care of PLWHA. Several risk factors for depressive symptoms among PLWHA receiving ART have been reported. Female patients have been reported to be at increased risk for depressive symptoms in most studies [24, 25]. A complex interaction between age and depressive symptoms has been identified [26]. Studies assessing residential status have also shown inconsistent results [27]. However, most previous studies were undertaken in high-income countries, and relatively few studies focused on this issue in low-and-middle-income countries, including Nigeria [28]. In this study, we aimed to explore the prevalence of and risk factors for depressive symptoms among PLWHA receiving ART in Nigeria to narrow this gap in literature, and provide background information for holistic health care for PLWHA.

## Method

### Study design and setting

This study employed a cross-sectional quantitative survey research design using a structured, interviewer administered clinical assessment tool. The study was carried out in three designated clinical hospitals in Birnin Kebbi, and Arungu of Kebbi State, Nigeria. Nigeria is made up of 36 states and the FCT, Abuja which are further grouped into six geopolitical zones (GPZs). Three GPZs are located in northern Nigeria i.e. North-east, North-central, and North-west GPZs, while southern Nigeria comprises the South-west, South-east and South-south GPZs. Kebbi is part of the North-west GPZ. This study was conducted in three HIV treatment centres in Kebbi. HIV treatment centres are health centres dedicated to provide antiretroviral therapy (ART) to PLWHAS. These

centres are run by government and non-governmental organisations in Nigeria.

### Study population and sample

The study was carried out among the PLWHAS who were receiving services at HIV treatment centres located in the Argungu and Birnin Kebbi, Kebbi State as at November 2019. Based on available records from the National Agency for the Control of AIDS (NACA) in Nigeria, there was an estimated population of 938,443 PLWHA who were either in-care or on ART at 18 dedicated treatment centres in Kebbi State. The study sample was drawn from three conveniently sampled HIV treatment centres located in the Kebbi State. These treatment sites were selected based mainly on the availability of well-kept patient's records which facilitated the randomization of the study participants and the ability to retrieve contact details of potential study participants. The recruitment criteria were as follows: (1) aged over 18 years and above, (2) confirmed diagnosis of HIV infection, (3) confirmed ART in Kebbi State, and (4) available to participate in the survey. The individual inclusion criteria for participating in this study seeking or receiving treatment in one of the three HIV treatment centres involved in this study. The exclusion criteria were: being under the age of 18, and not being able to communicate in English or Hausa languages.

### Measures

Socio-demographic questionnaire created by the research team, epidemiological and clinical data were directly extracted from the case reporting database of the China HIV/AIDS Information Network. Depressive symptoms was measured using the validated Beck Depression Inventory (BDI), a self-rated depression screening instrument with 21 items. It has been applied to assess the presence and severity of depressive symptoms in those greater than 13 years old [29], and has been reported to be an adequate measure for assessing depressive symptoms among PLWHA [30, 31]. Higher scores suggest greater depressive symptoms, with scores  $\geq 14$  and  $\leq 19$  reflecting mild depressive symptoms and scores  $\geq 20$  reflecting moderate to severe depressive symptoms.

Paper-based BDI questionnaires were given to PLWHA during their routine medical visit to the designated clinical hospitals during the study period. When informed consents were obtained, participants were required to fill out questionnaires in a private and quiet room under the supervision of a professional clinician prior to their regular physical examination and consultation. Every investigation lasted nearly half an hour.

### Statistical analyses

Descriptive statistics were used to summarize participants' characteristics. Means and standard deviations were used for continuous variables and frequencies and proportions were used for categorical variables. Multinomial regression analysis was used to explore the risk factors for depressive symptoms, with results presented by odds ratio (OR) and 95% confidence interval (CI).  $P < 0.05$  was considered significant statistically.

## Results

### Demographic details of Participants

Participants in the study came from northwest GPZs which

is one of the GPZs in Nigeria. A total of 242 (69.5%) and 106 (30.5%) participants were interviewed in English and Hausa languages, respectively. The sample comprised 24.4% female (n = 85) and 75.6% (n = 263) males. Table 1

shows the main demographic and clinical characteristics with significant differences among participants with minimal, mild and moderate to severe depressive symptoms.

**Table 1:** The comparison of characteristic among participants with different level of depressive symptoms

	Minimal	Mild	Moderate to severe	Statistics <sup>a</sup>	P value
BDI scores	5.78 ± 3.65	15.25 ± 1.61	30.27 ± 7.67	792.202	<0.002
Age/year	34.65 ± 12.14	37.86 ± 14.52	40.83 ± 14.07	4.407	0.013
<b>Gender</b>					
Male	122(42.6%)	58(22.1%)	83(31.5%)	3.234	0.312
Female	43(50.6%)	15(17.6%)	27(31.7%)		
<b>Marital status</b>					
Married/living with a partner	126(67.7%)	29(15.6%)	31(16.6%)	8.163	0.007
Unmarried/divorced/widowed	89(63.2%)	33(11.0%)	77(25.8%)		
<b>Residence</b>					
Urban	155(59.4%)	29(11.1%)	77(29.5%)	7.165	0.042
Rural	82(50.6%)	35(21.6.8%)	45(27.7%)		
<b>History of HCV infection</b>					
No	119(47.0%)	45(17.7%)	89(35.2%)	10.343	0.028
Yes	4(23.5%)	8(47.0%)	5(29.4%)		
Unknown	28(35.9%)	25(32.0%)	25(32.0%)		
<b>Recent Experience of side effects</b>					
Yes	221(61.8%)	28(11.2%)	37(27.1%)	13.603	0.001
No	32(51.6%)	16(25.8%)	14(22.5%)		

Depression Scale; HCV: Hepatitis C virus

<sup>a</sup>Chi-square test were used for categorical variables, one-way ANOVA were used for continuous variables

**Table 2:** Multinomial regression analysis for exploring risk factors for depressive symptoms

	Mild		Moderate to Severe	
	OR (95% CI)	P value	OR (95% CI)	P value
Age	1.03 (0.82, 1.08)	0.510	1.08 (1.09, 2.08)	0.032
<b>Gender</b>				
Female	Reference	Reference		
Male	2.20 (1.14, 6.11)	0.719	0.74 (0.55, 1.78)	0.382
<b>Residence</b>				
Rural	Reference	Reference		
Urban	0.66 (0.59, 1.84)	0.881	1.37 (0.75, 2.42)	0.151
<b>Marital status</b>				
Unmarried	Reference	Reference		
Married	2.20 (1.06, 5.32)	0.033	1.13 (0.52, 2.12)	0.562
<b>Recent experience of side effects</b>				
No	Reference	Reference		
Yes	2.38 (1.35, 5.82)	0.003	1.11 (0.43, 2.72)	0.812
<b>History of HCV infection</b>				
No	Reference	Reference		
Yes	4.19(1.21, 14.93)	0.021	1.89 (0.52, 6.41)	0.298
Unknown	0.65(0.21, 2.78)	0.631	1.74 (0.73, 4.77)	0.202

Abbreviations: OR: Odds Ratio; CI: Confidence Interval a Minimal depressive symptoms was created as a dummy variable and adjusted for age, gender and marital status

**Regression analysis**

Table 2 shows that mild depressive symptoms were significantly associated with being married or living with a partner (OR = 2.20, 95% CI, 1.14–6.11), experience of side effects (OR = 2.38, 95% CI, 1.35–5.82), and history of HCV infection (OR = 4.19, 95% CI, 1.21–14.93), while moderate to severe depressive symptoms were significantly associated with increasing age (OR = 1.08, 95% CI, 1.09–2.08).

**Discussion**

The study found that the prevalence of depressive symptoms among PLWHA receiving ART was 41.2 per cent, with 12.9 per cent having mild depressive symptoms and 25.3 per cent

having moderate to severe depressive symptoms. Being married or living with a partner, experiencing ART-related side effects and a history of HCV infection, independent risk factors for mild depressive symptoms were established, while increasing age was an independent risk factor for moderate to severe depressive symptoms. Being married or living with a partner, recent experience of side effects, history of HCV infection has been associated with 2.20, 2.38, 4.19 times the risk of mild depressive symptoms. Meanwhile, each one year increase in age was found to cause an approximately 2% increase in the odds of moderate to severe depressive symptoms. Our findings were supported by other relevant researches in varied populations, such as Korea, Australia, USA and Spanish [32-35]. Nevertheless, the risk factors for mild and moderate to severe depressive symptoms were different. In patients with mild to severe depressive symptoms, rising age played a more significant role than marital status, recent experience with side effects and history of HCV infection. It could probably be clarified that depressive symptoms were generally ignored in PLWHA [36], and that age could increase the progression of ignored depressive symptoms to more severe status due to age-related risk factors such as social isolation [37]. For patients with mild depressive symptoms, marital status, recent experience with side effects and history of HCV infection have played a more significant role than age. It could probably be clarified that, over time, mild depressive symptoms could be relieved by partner social support [38], treatment and alleviation of side effects [39] and HCV infection [40].

**Limitations**

Not all prospective participants who were originally routinely screened for this study could be contacted due to out-of-date contact information in clinical records. Therefore, several participants were contacted during their clinical days. There was also a difference between the real



research group in which the participants fall and the list of participants. This study had limitations. First, what we found was based on cross-sectional evidence, the causal relationship needs to be explained. Furthermore, some previously reported risk factors, such as HIV stigma, inadequate social support, income and alcohol use, have not been addressed in this review, which has made our results potentially biased. It might be best to consider these potential risk factors in any related investigations. However, we found a correlation between marital status, side effects of ART, history of HCV infection, increased age and depressive symptoms.

### Conclusion

In conclusion, marital status, ART-related side effects, history of HCV infection and aging were positively associated with depressive symptoms in PLWHA receiving ART. It suggested the need for urgent psychological care for PLWHA in ART, particularly for those of growing age, married or living with a partner, experience with side effects, and history of HCV infection.

### Reference

- Lopez DA, Mathers DC, Ezzati M, Jamison TD, Murray JLC. Global burden of disease and risk factors. New York: Oxford University Press/The World Bank, 2006.
- Fortes S, Villano LAB, Lopes CS. Nosological profile and prevalence of common mental disorders of patients seen at the Family Health Program (FHP) units in Petrópolis, Rio de Janeiro. *Rev Bras Psiquiat* 2008;30(1):32-7.
- Chibanda D, Cowan F, Gibson L, Weiss HA, Lund C. Prevalence and correlates of probable common mental disorders in a population with high prevalence of HIV in Zimbabwe. *BMC Psychiatry* 2016;16(1):55.
- Primeau MM, Avellaneda V, Musselman D, St Jean G, Illa L. Treatment of depression in individuals living with HIV/AIDS. *Psychosomatics* 2013;54(4):336-44.
- Kidia K, Machando D, Bere T, Macpherson K, Nyamayaro P, Potter L *et al.* 'I was thinking too much': experiences of HIV-positive adults with common mental disorders and poor adherence to antiretroviral therapy in Zimbabwe. *Tropical Medicine & International Health* 2015;20(7):903-13.
- Yi S, Chhoun P, Suong S, Thin K, Brody C, Tuot S. AIDS-related stigma and mental disorders among people living with HIV: a cross-sectional study in Cambodia. *PLoS One* 2015;10(3):e0121461.
- Nakimuli-Mpungu E, Bass JK, Alexandre P, Mills EJ, Musisi S, Ram M, Katabira E. Depression, alcohol use and adherence to antiretroviral therapy in sub-Saharan Africa: a systematic review. *AIDS and Behavior* 2012;16(8):2101-18.
- Igie I, Udoh M: Prevalence of depression in HIV/AIDS patients, Lagos state university hospital (LUTH), Lagos, Nigeria. *International AIDS society*. In: 6th IAS conference on HIV Pathogenesis, Treatment and Prevention: Rome, Italy 2011.
- Adewuya AO, Afolabi MO, Ola BA, Ogundele OA, Ajibare AO, Oladipo BF. Relationship between depression and quality of life in persons with HIV infection in Nigeria. *The International Journal of Psychiatry in Medicine* 2008;38(1):43-51.
- Olisah V, Baiyewu O, Sheikh T. Adherence to highly active antiretroviral therapy in depressed patients with HIV/AIDS attending a Nigerian university teaching hospital clinic: original article. *Afr J Psychiatry* 2010;13(4):275-9
- Sale S, Gadanya M. Prevalence and factors associated with depression in HIV/AIDS patients aged 15–25 years at Aminu Kano teaching hospital, Nigeria. *J Child and Adolesc Mental Health* 2008;20(2):95-9
- Ndu A, Arinze-Onyia S, Aguwa E, Obi I. Prevalence of depression and role of support groups in its management: a study of adult HIV/AIDS patients attending HIV/AIDS clinic in a tertiary health facility in South-eastern Nigeria. *J Public Health Epidemiol.* 2011;3(4):182-6.
- Global Health Observatory (GHO) data - HIV/AIDS [<http://www.who.int/gho/hiv/en/>.]
- Nigeria: WHO statistical profile [<http://www.who.int/gho/countries/nga.pdf?ua=1>.]
- HIV and AIDS in Nigeria [<http://www.avert.org/professionals/hiv-aroundworld/sub-saharan-africa/nigeria>.]
- Whiteside A. HIV/AIDS: a very short introduction. Oxford: Oxford University Press 2008.
- Oguntibeju OO. Quality of life of people living with HIV and AIDS and antiretroviral therapy. *HIV/Aids (Auckland, NZ)* 2012;4:117.
- Jiang Y, Wang M, Wei XQ, He JM, Guo TS, Huang GQ. Prevalence of depression and related factors in 180 HIV/AIDS patients receiving highly active antiretroviral therapy. *Zhonghua liu Xing Bing xue za zhi= Zhonghua Liuxingbingxue Zazhi* 2016;37(5):638-42.
- Niu L, Luo D, Liu Y, Silenzio VM, Xiao S. The mental health of people living with HIV in China, 1998–2014: a systematic review. *PLoS One* 2016;11(4):e0153489.
- Gu L, Xie J, Long J, Chen Q, Chen Q, Pan R *et al.* Epidemiology of major depressive disorder in mainland china: a systematic review. *PLoS one* 2013;8(6):e65356.
- Rivera-Rivera Y, Vázquez-Santiago FJ, Albino E, Sánchez MD, Rivera-Amill V. Impact of depression and inflammation on the progression of HIV disease. *Journal of clinical & cellular immunology*, 2016, 7(3).
- Sherr L, Clucas C, Harding R, Sibley E, Catalan J. HIV and depression—a systematic review of interventions. *Psychology, health & medicine* 2011;16(5):493-527.
- Wagner GJ, Ghosh-Dastidar B, Robinson E, Ngo VK, Glick P, Mukasa B. Effects of depression alleviation on ART adherence and HIV clinic attendance in Uganda, and the mediating roles of self-efficacy and motivation. *AIDS and Behavior* 2017;21(6):1655-64.
- Brandt R. The mental health of people living with HIV/AIDS in Africa: a systematic review. *African Journal of AIDS Research* 2009;8(2):123-33.
- Robertson K, Bayon C, Molina JM, McNamara P, Resch C, Muñoz-Moreno JA *et al.* Screening for neurocognitive impairment, depression, and anxiety in HIV-infected patients in Western Europe and Canada. *AIDS care* 2014;26(12):1555-61.
- Nanni MG, Caruso R, Mitchell AJ, Meggiolaro E, Grassi L. Depression in HIV infected patients: a review. *Current psychiatry reports* 2015;17(1):530.
- Rodriguez V, Magis C, Casillas J, Ramirez HV, Hernandez G, Rentería CR. Internal consistency and

- factorial structure of the Beck Depression Inventory (BDI) in Mexican HIV-positive patients. InPoster at XIX International AIDS Conference 2012,22-27.
28. Li XK, Ji WJ, Zhao J, Wang SJ, Au CT. Ammonia decomposition over Ru and Ni catalysts supported on fumed SiO<sub>2</sub>, MCM-41, and SBA-15. *Journal of Catalysis* 2005;236(2):181-9.
  29. Rong H, Nianhua X, Jun X, Lianguo R, Si W, Sheng W. Prevalence of and risk factors for depressive symptoms among people living with HIV/AIDS receiving antiretroviral treatment in Wuhan, China: a short report. *AIDS care* 2017;29(12):1524-8.
  30. Mohammed A, Said JM, Wakil MA, Rabbebe IB, Sheikh T, Agunbiade S. Unrecognized psychiatric disorders among adult patients admitted into a general hospital in Maiduguri, Northeastern Nigeria. *Pan Afr Med J* 2014;19:197.
  31. Hobkirk AL, Starosta AJ, De Leo JA, Marra CM, Heaton RK, Earleywine M. Psychometric validation of the BDI-II among HIV-positive CHARTER study participants. *Psychological assessment* 2015;27(2):457.
  32. Kee MK, Lee SY, Kim NY, Lee JS, Kim JM, Choi JY *et al.* Anxiety and depressive symptoms among patients infected with human immunodeficiency virus in South Korea. *AIDS care* 2015;27(9):1174-82.
  33. Kim MH, Mazenga AC, Yu X, Devandra A, Nguyen C, Ahmed S. Factors associated with depression among adolescents living with HIV in Malawi. *BMC psychiatry* 2015;15(1):264.
  34. Libman H, Saitz R, Nunes D, Cheng DM, Richardson JM, Vidaver J *et al.* Infection is associated with depressive symptoms in HIV-infected adults with alcohol problems. *The American journal of gastroenterology* 2006;101(8):1804.
  35. Pakianathan M, Whittaker W, Lee MJ, Avery J, Green S, Nathan B. Chemsex and new HIV diagnosis in gay, bisexual and other men who have sex with men attending sexual health clinics. *HIV medicine* 2018;19(7):485-90.
  36. Cholera R, Pence BW, Bengtson AM, Crane HM, Christopoulos K, Cole SR *et al.* Mind the gap: gaps in antidepressant treatment, treatment adjustments, and outcomes among patients in routine HIV care in a multisite US Clinical Cohort. *PloS one* 2017;12(1):e0166435.
  37. Rong H, Nianhua X, Jun X, Lianguo R, Si W, Sheng W. Prevalence of and risk factors for depressive symptoms among people living with HIV/AIDS receiving antiretroviral treatment in Wuhan, China: a short report. *AIDS care* 2017;29(12):1524-8.
  38. Umeadi CA. Effect of Social Support and HIV-Related Stigma on Depression in HIV/AIDS Patients.
  39. Chen WT. Side effects of antiretroviral therapy (ART) are associated with depression in Chinese individuals with HIV: a mixed methods study. *Journal of midwifery & women's health* 2013;58(5):585.
  40. Feuillet P, Lert F, Tron L, Aubriere C, Spire B, Dray-Spira R. Agence Nationale de Recherche sur le Sida et les Hépatites Virales (ANRS)-VIH: Enquête sur les personnes atteintes (Vespa2) Study Group. Prevalence of and factors associated with depression among people living with HIV in France. *HIV Med*, 2016.